

By inhibiting NLRP3 inflammasome via autophagy activation, carnosol improves sevoflurane-induced cognitive impairment in old rats.

**Dr.V.Ravikumar¹, Dr.P.Aravinda Reddy², Shilpadas³, Ashaful alom⁴,
Assistant professor^{1,2,3,4},
Department of Pharmacy,
Samskruti College of Pharmacy,
Kondapur (V), Ghatkesar (M) Medchal Dist, Telangana, India.**

Abstract:

Specifically, we want to learn how carnosol works and how it affects postoperative cognitive dysfunction (POCD). Methods: The sevoflurane (SEV) paradigm of cognitive impairment in rats was developed. We used a dihydroethidium (DHE) test to measure the impact of carnosol on ROS levels in rats that had been produced by sevoflurane. The impact of carnosol on mitochondrial damage was evaluated using adenosine triphosphate (ATP) generation and immunoblot tests. Cognitive impairment in the rats was assessed by a water maze experiment. The immunoblot test was used to evaluate the action mechanism. The results showed that carnosol inhibited ROS generation and mitochondrial damage caused by sevoflurane in rats, and it also triggered autophagy. Carnosol also inhibited SEV-induced cognitive impairment via activating autophagy and SEV-induced NLRP3 inflammasome activation, respectively. Conclusion: Carnosol, by inhibiting the NLRP3 inflammasome, improves SEV-stimulated cognitive impairment and may one day be used as a treatment for cognitive impairment. Carnosol, Sevoflurane (SEV), NLRP3 inflammasome, and autophagy are some of the terms used to describe this phenomenon.

INTRODUCTION

Postoperative cognitive dysfunction (POCD) is a type of complication with long-term consequences, defined as impaired memory, attention and information processing that occurs after anesthesia [1]. Studies show that about 10 % of surgery patients as well as 40 % of older patients over the age of 65 develop POCD [2]. Compared to patients without POCD, patients with POCD have significantly higher mortality rates and higher dependence on social security [3]. Despite extensive research efforts, the pathogenesis of POCD remains unknown [4]. New therapeutic agents still need to be developed to treat POCD. Rosemary (*Salvia rosmarinus*) is a Mediterranean plant that is now widely available in several countries [5,6]. Rosemary extract can be used to treat inflammation-related diseases [7]. Carnosol is a phenolic diterpenoid which exists in rosemary extract and has been shown to have anti-

inflammatory and antioxidant effects [8]. Carnosol treatment inhibited the eosinophils in the bronchoalveolar lavage fluid of mice after ovalbumin treatment [9]. Previous studies have shown that carnosol alleviates sevofluranestimulated cognitive dysfunction in aged rats via NF- κ B pathway. It also causes the activation of autophagy in breast cancer [8]. However, the potential effect and mechanism of action of the compound on POCD remains unknown. There is growing evidence of a causal relationship between SEV-stimulated cognitive impairment and NLRP3 inflammasome [10]. In response to cellular stress, NLRP3 recruits ASC as well as pro-caspase-1, which causes the cleaved caspase-1 to activate and process the maturation of IL-1 β and IL-18 [11]. NF- κ B is involved in the control of many cellular processes [11]. The activation of NF- κ B family is a key step in regulating pyroptosis. However,

the effects of carnosol on autophagy and its relationship with NLRP3 is unclear. The aim of this study was to investigate the role of carnosol in POCD progression using a sevoflurane-induced cognitive impairment rat model of rats.

EXPERIMENTAL

POCD rat model Male Sprague–

Dawley rats (12 months old, 250 – 280 g) were purchased from Vital River (Beijing, China). The procedures in this study were approved by the Ethics Committee of Xi'an International Medical Center Hospital, Shanxi, China (approval no. 202315), and conducted in accordance with the guidelines of Declaration of Helsinki in Biomedical and Behavioral Research [12]. For the establishment of POCD model, sevoflurane (2 % SEV) was inhaled by old rats (8 months old) for 5 h, and carnosol administered to the rats in doses of 5 and 10 mg/kg every 2 days, respectively. The rats were divided into the following groups: control, SEV, SEV + carnosol (5 mg/kg), as well as SEV + carnosol (10 mg/kg) groups.

Morris water maze (MWM) tests

The MWM tests were performed in a swimming pool with four quadrants and the activity of the rats was recorded. The rats were trained for 4 days in order to determine the location of the platform; if the rats could not find the location within 90 s, they were placed on the platform. Finally, the platform was removed. The time and number was recorded.

Determination of ROS

For ROS staining, the cells were mounted and stained with Dihydroethidium (DHE). DHE fluorescence of 20 cells per field in five random fields was quantified by the use of AxioVision software.

JC-1 staining

To analyze mitochondria damage, samples were placed in 12-well plates and kept for 24 hours for cell adhesion. After rinsing, the cells were

incubated with 2 μ M JC-1 for 15 min at 37 °C. After rinsing in phosphate buffer (PBS) 3 times, the cells were photographed and the JC-1 positive cells were counted per field.

Determination of ATP production

For the determination of ATP production, an ATP detection kit (Beyotime, Beijing, China) was utilized in accordance with the manufacturer's protocol.

Immunoblot assay

The samples were electrophoresed and transferred onto PVDF membranes and then blocked with 5 % fat-free milk. Subsequently, the membranes were conjugated with antibodies targeting LC3 (ab192890, 1:1000, Abcam), p62 (ab109012, 1:1000, Abcam), NLRP3 (ab263899, 1:1000, Abcam), ASC (ab283684, 1:500, Abcam), Caspase-1 (ab207802, 1:500, Abcam), IL-18 (ab243091, 1:1000, Abcam), IL-1 β (ab254360, 1:1000, Abcam), and β -actin (ab8226, 1:3000, Abcam) for 2 h at room temperature. Subsequently the membranes were incubated with secondary antibodies (1:2000, Abcam) for 1 h at room temperature. The blots were then analyzed with ECL kit.

Statistics

Statistical analysis was performed with GraphPad 7.0 software using t-test, and the data are shown as mean \pm SD. $P < 0.05$ was considered statistically significant.

RESULTS

Carnosol suppressed sevoflurane-induced ROS production of rats. The results showed that carnosol treatment decreased DHE expression levels in sevoflurane-induced hippocampal neurons, suggesting the inhibition of ROS production (Figure 1). Therefore, Carnosol suppressed sevoflurane-induced ROS production of rats.




Figure 1: Carnosol suppressed sevoflurane induced ROS production of rats. DHE was used to detect ROS levels in hippocampus from control, SEV, SEV+carnosol (5 mg/kg), and SEV+carnosol (10 mg/kg) groups. Scale bar indicates 100 μ m

Carnosol suppressed mitochondrial damage induced by sevoflurane, and activated autophagy

Immunostaining showed the degree of mitochondrial damage of the hippocampus from each group. The aggregates and monomers reflected the degree of mitochondrial damage. The data showed that the monomers were enhanced and the aggregates decreased in SEV hippocampal neurons (Figure 2 A). In addition, carnosol reduced the monomers and increased the aggregates in SEV-induced hippocampal neurons, suggesting the inhibition of mitochondrial damage (Figure 2 A). Similarly, carnosol decreased ATP production in SEV hippocampal neurons but induced the production of ATP in SEV hippocampal neurons (Figure 2 B). Western blot results showed that the ratio of LC3BII/LC3BI in SEV hippocampal neurons increased, while carnosol decreased the ratio in SEV hippocampal neurons, suggesting the inhibition of autophagy (Figure 2 C). Interestingly, increased expression of p62 was found in SEV hippocampal neurons but carnosol decreased p62 expression in SEV hippocampal neurons, further confirming the previous data (Figure 2 D). Therefore, carnosol alleviated mitochondrial damage induced by sevoflurane and activated autophagy.




Figure 2: Carnosol suppressed mitochondrial damage induced by sevoflurane and activates autophagy. (A). Immunostaining showed the degree of mitochondrial damage of hippocampus from the indicated groups. Green indicates monomers; Red indicates aggregates. Scale bar indicates 50 μ m; (B). ATP production was shown in hippocampus from the indicated groups; (C). Immunoblot assays showed the expression levels of LC3B in hippocampus from the indicated groups. The ratio of LC3BII/LC3BI was calculated; (D). Immunoblot assays showed the expression levels of p62 in hippocampus from the indicated groups. Data are presented as mean \pm SD; ***P < 0.001, SEV vs control, ##p < 0.01, ###p < 0.001, SEV+carnosol vs SEV

Carnosol suppressed sevoflurane-induced NLRP3 inflammasome activation

To determine the potential mechanism of carnosol-mediated effect in sevoflurane induced rats, the role of carnosol in the NLRP3 pathway was assessed after treatment with SEV, carnosol and the inhibitor of autophagy, 3-MA. Carnosol increased the levels of NLRP3 in SEV group was found, and reversed the increase of NLRP3 expression caused by SEV treatment (Figure 3 A). However, 3-MA treatment further increased the expression of NLRP3 in carnosol-treated rats, suggesting the regulation of autophagy in NLRP3 pathway (Figure 3 A). Furthermore, NLRP3 signaling pathway was activated following the establishment of SEV-induced model as demonstrated by increased levels of

ASC, IL-1 β and IL-18 (Figure 3 B). The decreased levels of ASC, IL-1 β and IL-18 levels were caused by carnosol treatment (Figure 3 B). 3-MA treatment increased further the expression of NLRP3 in carnosol-treated rats. Thus, the data suggest that carnosol suppressed sevoflurane-induced NLRP3 inflammasome activation by regulating autophagy.




Figure 3: Carnosol suppressed sevoflurane induced NLRP3 inflammasome activation by activating autophagy. (A). Immunoblot assays showed the expression levels of NLRP3 in hippocampus from the indicated groups; (B). Immunoblot assays showed the expression levels of indicated proteins in the hippocampus from the indicated groups. Data are presented as mean \pm SD, ***P < 0.001, SEV vs control, ###p < 0.001, SEV+carnosol vs SEV, \$\$\$p < 0.001, SEV+carnosol+3-MA vs SEV+carnosol




Figure 4: Carnosol suppressed sevoflurane induced cognitive dysfunction by inhibiting

NLRP3 inflammasome activation. (A). Water maze experiment reveals rat movement track from the indicated groups; (B-E). Escape latency (B), time of across the stealth platform (C), time in target quadrant (D), and distance in target quadrant (E) of rats from the indicated groups. Data are presented as mean \pm SD; ***p < 0.001, SEV vs control, #p < 0.05, ##p < 0.01, ###p < 0.001, SEV+carnosol vs SEV, \$p < 0.05, \$\$\$p < 0.001, SEV+carnosol+3-MA vs SEV+carnosol

Carnosol suppresses sevoflurane-induced cognitive dysfunction by inhibiting NLRP3 inflammasome activation

As expected, the cognitive ability of rats in SEV group was impaired significantly with poor ability to escape latency, locate the target, long latency time and longer time in the target quadrate (Figure 4 A - E). Carnosol treatment improved the performance of rats (Figure 4 B - E). Interestingly, NLRP3 treatment further impaired the performance of rats, suggesting that carnosol alleviated cognitive dysfunction (Figure 4 B - E). Therefore, carnosol suppressed SEV-induced cognitive dysfunction by suppressing NLRP3 inflammasome activation.

DISCUSSION

With the different types of surgery and the duration of general anesthesia, the degree of POCD varies. In recent years, there has been great progress in anesthesia technology, monitoring instruments and surgical operations, and these have greatly improved the safety of surgery, POCD is still common and its pathogenesis and mechanisms have attracted wide attention [10]. POCD can usually be treated with medication [10]. If the patient has cognitive dysfunction after surgery, it can be treated with drugs, as the main purpose is to improve the patient's condition and facilitate recovery [10]. With the aid of intravenous drip, nutritional neurodrugs and other treatments to ameliorate brain tissue ischemia and hypoxia, and postoperative recovery, using mecobalamine tablets, adenosine monophosphate and other drugs, may improve the cognitive ability of patients [13]. However, it is still necessary to find more effective drugs. Here, carnosol was

shown to ameliorate the sevoflurane-induced cognitive impairment in aged rats. Therefore, carnosol may serve as a drug for the management of POCD.

Carnosol is a diterpenoid phenolic compound, which has antioxidant, anti-inflammatory, antiproliferation and anti-tumor effects [14,15]. Due to its remarkable pharmacological activity, more and more scholars at home and abroad have paid more attention to it in recent years [8]. It has a potential role in the prevention and treatment of neuropathic diseases [8]. Carnosol is a potent ribosome S6 kinase (RSK2) inhibitor and also a carnosol is also an Nrf2 activator, increasing Nrf2 levels and promoting heme oxygenase-1 (HMOX1) expression [16]. It alleviated mitochondrial damage induced by SEV and also activated autophagy and increased SEV-induced NLRP3 inflammasome activation by activating autophagy. Carnosol ameliorated sevoflurane-induced cognitive impairment. Previous studies have shown that it alleviated sevoflurane-induced cognitive dysfunction in aged rats by mediating NF- κ B pathway [16].

In the present study, carnosol alleviated sevoflurane-induced cognitive dysfunction by regulating NLRP3 inflammasome as well as autophagy. However, further in vivo studies to elucidate the mechanism of action are still needed. Autophagy is vital in the elimination of protein aggregates and abnormal organelles [10]. Enhanced autophagy disrupts neuronal homeostasis [17]. Activation of autophagy suppresses the formation of NLRP3 inflammasome [10]. NLRP3 inflammasome is highly expressed and over-activated in aged rats, and is vital in the central inflammatory response [18]. Over-activation of NLRP3 inflammasome leads to a variety of diseases, including POCD [10]. NLRP3 inflammasomes are involved in the important mechanisms of POCD [10]. It also reduced cognitive impairment in older rats. Interestingly, carnosol further improved SEV-induced NLRP3 inflammasome activation by activating autophagy, thus ameliorating POCD. However, the precise mechanism of action needs to be investigated further.

CONCLUSION

Carnosol ameliorates SEV-induced cognitive impairment in aged rats by suppressing NLRP3 inflammasome, as well as autophagy. Therefore, carnosol is a promising drug for the management of POCD.

REFERENCES

1. Li X, Gao Y, Han X, Tang S, Li N, Liu X, Ni X. *Maresin1 ameliorates postoperative cognitive dysfunction in aged rats by potentially regulating the NF-kappaB pathway to inhibit astrocyte activation. Exp Gerontol 2023; 176: 112168.*
2. Wang G, Shen J, Zhai L, Lin Y, Guan Q, Shen H. *TL1A promotes the postoperative cognitive dysfunction in mice through NLRP3-mediated A1 differentiation of astrocytes. CNS Neurosci Ther 2023; 29(11):3588-3597.*
3. Wang X, Chen X, Wu F, Liu Y, Yang Y, Chen W, Pan Z, Hu W, Zheng F, He H. *Relationship between postoperative biomarkers of neuronal injury and postoperative cognitive dysfunction: A meta-analysis. PLoS One 2023; 18(4): e0284728.*
4. Zhao W, Li J, Wang J, Liu M, Yu D. *Effect of dexmedetomidine on postoperative cognitive dysfunction and the T helper 17/regulatory T cell balance in geriatric patients undergoing orthopedic surgery: a randomized controlled study. Am J Transl Res 2023; 15(4): 2634-2644.*
5. Farouk SM, Abdel-Rahman HG, Abdallah OA, El-Behidy NG. *Comparative immunomodulatory efficacy of rosemary and fenugreek against Escherichia coli infection via suppression of inflammation and oxidative stress in broilers. Environ Sci Pollut Res Int 2022; 29(26): 40053-40067.*
6. Ma W, Shi H, Wei G, Hua M, Yu H. *Loureirin B attenuates amiodarone-induced pulmonary fibrosis by suppression of TGF β 1/Smad2/3*

- pathway. *Trop J Pharm Res* 2020; 19(7): 1371-1376.
7. Bai R, Yuan C, Wang T, Liu L, Li J, Lai Y, Li H, Chen Z, Li C, Ke D et al. Apple pomace and rosemary extract ameliorates hepatic steatosis in fructose-fed rats: Association with enhancing fatty acid oxidation and suppressing inflammation. *Exp Ther Med* 2020; 20(3): 1975-1986.
 8. Gonzalez-Cardenete MA, Gonzalez-Zapata N, Boyd L, Rivas F. Discovery of Novel Bioactive Tanshinones and Carnosol Analogues against Breast Cancer. *Cancers (Basel)* 2023; 15(4): 1318.
 9. Lee JE, Im DS. Suppressive Effect of Carnosol on Ovalbumin-Induced Allergic Asthma. *Biomol Ther (Seoul)* 2021; 29(1): 58-63.
 10. Zhou J, Zhang C, Fang X, Zhang N, Zhang X, Zhu Z. Activation of autophagy inhibits the activation of NLRP3 inflammasome and alleviates sevoflurane-induced cognitive dysfunction in elderly rats. *BMC Neurosci* 2023; 24(1): 9.
 11. Han QQ, Le W. NLRP3 Inflammasome-Mediated Neuroinflammation and Related Mitochondrial Impairment in Parkinson's Disease. *Neurosci Bull* 2023; 39(5): 832-844.
 12. World Medical A. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *J Postgrad Med* 2002; 48(3): 206-208.
 13. Wang W, Zhao B, Gao W, Song W, Hou J, Zhang L, Xia Z. Inhibition of PINK1-Mediated Mitophagy Contributes to Postoperative Cognitive Dysfunction through Activation of Caspase-3/GSDME-Dependent Pyroptosis. *ACS Chem Neurosci* 2023; 14(7): 1249-1260.
 14. Habtemariam S. Anti-Inflammatory Therapeutic Mechanisms of Natural Products: Insight from Rosemary Diterpenes, Carnosic Acid and Carnosol. *Biomedicines* 2023; 11(2): 545.
 15. Luan Y, Jiang L, Luan Y, Xie Y, Yang Y, Ren KD. Mitophagy and Traumatic Brain Injury: Regulatory Mechanisms and Therapeutic Potentials. *Oxid Med Cell Longev* 2023; 2023: 1649842.
 16. Li X, Zhang Q, Hou N, Li J, Liu M, Peng S, Zhang Y, Luo Y, Zhao B, Wang S et al. Carnosol as a Nrf2 Activator Improves Endothelial Barrier Function Through Antioxidative Mechanisms. *Int J Mol Sci* 2019; 20(4): 880.
 17. Pajarillo E, Kim SH, Digman A, Dutton M, Son DS, Aschner M, Lee E. The role of microglial LRRK2 in manganese-induced inflammatory neurotoxicity via NLRP3 inflammasome and RAB10-mediated autophagy dysfunction. *J Biol Chem* 2023; 299(7): 104879.
 18. Wu Y, Hu A, Shu X, Huang W, Zhang R, Xu Y, Yang C. *Lactobacillus plantarum* postbiotics trigger AMPKdependent autophagy to suppress *Salmonella* intracellular infection and NLRP3 inflammasome activation. *J Cell Physiol* 2023; 238(6): 1336-1353.