

Oral Agarwood's Potential as a Treatment for Chronic Inflammatory Diseases

Dr. Priyanka Shrenik Gugale
Assistant Professor

Homoeopathic Medical College & Hospital, Jalgaon

Abstract: Gingivitis, in one form or another, may affect almost everyone. Many things, like not cleaning your teeth enough or an immune system that isn't working properly, may lead to gingivitis, a kind of periodontal disease. Periodontitis, which is irreversible, may develop from gingivitis if left untreated. Finding a medication that may decrease inflammation with fewer adverse effects, as an alternative to existing anti-inflammatory medicines, is a significant avenue of gingivitis research. Traditional Chinese medicine makes extensive use of agarwood for its therapeutic properties. According to prior research on agarwood's chemical composition and pharmacological actions, sesquiterpene and quercetin, two of the plant's active components, can reduce inflammation. Nevertheless, agarwood's potential in alleviating chronic inflammation in the mouth, including gingivitis, has received little scientific attention. This research aimed to outline the potential mechanism of action of agarwood on gingivitis by reviewing and synthesizing the relevant domestic and international literature. To determine how agarwood could work to reduce chronic inflammation in the mouth, we looked at the anti-inflammatory, antioxidant, antibacterial, analgesic, and immunological functions of its active ingredients in gingivitis. If you are interested in using agarwood or one of its byproducts to alleviate inflammation in your mouth, this review will serve as a useful theoretical reference.

Keywords: Agarwood, Gingivitis, Anti Inflammation, Bacteriostasis, Analgesia

Introduction

The resinous heartwood produced by trees belonging to the genus *Aquilaria* Lam is known as agarwood. According to taxonomy, it is a member of the *Daphne* family. *Aquilaria Sinensis*, scientifically known as *Aquilariasinensis* (Lou.) Gilg, is the primary component of agaricus grown in China [1]. Taiwan, Guangdong, Guangxi, and Hainan Island are the primary production regions [2]. Traditional medicine practitioners have relied on agarwood for centuries to alleviate a variety of symptoms, including heart problems, nausea, vomiting, and shortness of breath [3]. Traditional Chinese medicine also heavily relies on agarwood. It is anthelmintic and has a sedative effect. Stomach pain is another common usage for it.

issues, a runny nose, watery eyes, rheumatism, and a high temperature [4, 5]. The unique scent components of agarwood are refined and concentrated to create agarwood essential oil. According to studies, its primary components include chemicals that prevent the development of many illnesses, such as sesquiterpenes, aromatics, chromones, fatty acids, and others. Its many pharmacological effects have been carefully engineered [6]. Inflammation of the gums brought on by bacteria and tartar buildup in the gingival groove is known as gingivitis. Progression to periodontitis is possible if prompt treatment is not provided. Agarwood is a local specialty with great therapeutic potential and great scientific value. It would be wise to investigate its history, potential uses, and integration with diverse goods in a range of formats to fulfill its function.

1. Anti-Inflammatory Activity

Gingivitis is a chronic inflammatory disease, manifested as gingival swelling, bleeding and pain. Agarwood has anti-inflammatory effect, chromone and sesquiterpene are the main anti-inflammatory related active components isolated from agarwood essential oil [7]. The anti-inflammatory active components of agarwood can play an anti-inflammatory role by inhibiting the synthesis of inflammatory mediators such as 5-hydroxytryptamine, histamine and prostaglandin [8]. In molecular docking and ADME studies, it was found that several major sesquiterpenoids in agarwood essential oil have strong affinity for major anti-inflammatory receptors and immunoregulatory receptors [9]. Some sesquiterpenoid monomers also inhibit the release of NO induced by lipopolysaccharide in the anti inflammatory activity screening model of mouse monocyte macrophages RAW 264.7 in vitro [10, 11]. It was found that sesquiterpenoids in agarwood essential oil inhibit the expression of p-STAT3, thereby reducing the production of proinflammatory cytokine IL-1 β and IL-6, releasing anti-inflammatory mediators [12]. In addition, some scholars confirmed through experiments that 2

- (2-phenylethyl) chromone derivatives in agarwood can inhibit STAT1/3 and NF- κ B signal pathway showed significant anti-inflammatory effect [2]. Tryptone derivatives can also inhibit the production of carbon

monoxide in macrophages induced by lipopolysaccharide [1]. In addition, the alcohol extracts of the flowers, seeds, pericarp and leaves of agarwood contain n-hexadecanoic acid and squalene [1]. Hexadecanoic acid can inhibit phospholipase A2 by binding to the active site of phospholipase A2 to control inflammation [9]. Some researches have found that agarwood essential oil has significant anti-inflammatory activity when administered by gavage. The anti-inflammatory activity of agarwood essential oil has been confirmed. The incidence of periodontal tissue inflammation is high in systemic inflammation. At present, there is no precedent for the treatment of periodontitis with agarwood composition. The anti-inflammatory function of agarwood extract provides a theoretical basis for its application in the treatment of gingivitis.

2. Antioxidant Effects

A large number of studies have shown that oxidative stress is caused by the imbalance between the excessive production of reactive oxygen species (ROS) and the relative lack of antioxidants, and oxidative stress is also one of the pathophysiological mechanisms of periodontal tissue inflammation [13]. ROS can change the periodontal microenvironment by activating inflammatory factors, nuclear factor KB (NF-KB), c-Jun amino terminal kinases (JNKs) and autophagy, causing indirect serious damage to periodontal tissue, leading to periodontitis [14]. Some people measured the antioxidant activity of agarwood essential oil by 1,1-diphenyl-2-trinitrophenylhydrazine (DPPH) radical scavenging and iron ion reduction / antioxidant capacity method (FRAP), proving that it has a good antioxidant capacity in vitro [15]. Its essential oil extract β -Caryophyllene shows strong antioxidant effect [16]. Its antioxidation can reduce the level of intracellular reactive oxygen species to a certain extent, and significantly enhance the cellular antioxidation. The activity of enzyme SOD and GSH Px can protect PC12 cells from oxidative damage induced by H_2O_2 [17]. There is also a flavonoid compound in the agarwood extract that can provide antioxidant effect by preventing the formation of ROS, directly capturing ROS, protecting lipophilic antioxidants and stimulating the increase of enzymatic antioxidants [18]. In addition, different concentrations of agarwood essential oil have scavenging effects on DPPH free radicals, hydroxyl free radicals and superoxide anion free radicals, and with the increase of the concentration of agarwood essential oil, the antioxidant capacity gradually increases, with obvious correlation [19]. The analysis results also showed that both young leaves and mature leaves of agarwood have strong antioxidant activity, and the two kinds of leaves have the same use value in terms of chemical content and antioxidant capacity, which can reduce inflammatory reaction to a certain extent [20]. Agarwood essential oil shows certain antioxidant capacity in various systems, can reverse the imbalance between reactive oxygen species and antioxidants, and has the potential to be applied to gingivitis. Therefore, agarwood is a natural antioxidant, which can resist oxidative stress and alleviate gingivitis.

3. Antibacterial Activity

The antibacterial and anti-inflammatory effects of

agarwood are the main mechanisms to alleviate gingivitis. Gingivitis is caused by substances accumulated in or near the gingival sulcus by microbial plaque. Dental plaque is the main cause related to the development of gingivitis. It can aggravate inflammatory reaction by enhancing plaque accumulation or enhancing the susceptibility of gingival tissue to microbial attack. In recent years, it has been reported that agarwood has antibacterial activity and can inhibit some bacteria in oral cavity. Bacteria in the mouth mainly include *Streptococcus mutans*, *Porphyromonas gingivalis*, *Staphylococcus*, etc. The antibacterial activity test shows that agarwood essential oil has a good inhibitory effect on Gram-negative bacteria and Gram-positive bacteria, and has a more obvious antibacterial effect on Gram-positive bacteria. Because the cell wall of the external gram-negative bacteria contains a layer of lipopolysaccharide, which prevents hydrophobic compounds from entering the cells, thereby reducing the bacteriostatic effect [21]. In the process of agarwood formation, the infection of pathogenic bacteria causes the defense system of the *Aquilaria Sinensis* to resist external damage. It can prevent further infection of the pathogenic bacteria by generating local allergic reaction, that is, blocking the damage site [22]. This indicates that antibacterial activity has been carried out in the process of agarwood formation, so agarwood itself may have antibacterial activity. In previous studies, Mei showed that Chinese agarwood essential oil has anti MRSA activity [23]. Wetwitayaklung found that agarwood essential oil (*A. crassna*) has antibacterial activity against *Candida albicans* [24]. Filter paper method was used to test the antibacterial ability of agarwood essential oil by measuring the antibacterial circle. With the increase of the concentration of agarwood essential oil, the inhibition rate against bacteria increased. The inhibition rate of agarwood essential oil against bacteria is in the following order: *Staphylococcus aureus* > *Bacillus subtilis* > *Escherichia coli* [25]. The antibacterial effect of agarwood essential oil is related to sesquiterpenes and chromones in agarwood. Sesquiterpenes usually have antibacterial activity. Studies have found that sesquiterpenes in the volatile oil of agarwood β -Caryophyllene has significant antioxidant and antibacterial properties. Its antibacterial activity is attributed to its strong antioxidant activity [26]. Flavone widely exist in the plant kingdom, and are one of the main components of agarwood. The pharmacological activities of flavonoids include anti-inflammatory activity, anti allergic activity, antioxidant activity, anti-tumor activity, antifungal and antibacterial activity, which can inhibit a variety of bacterial virulence factors [27].

4. Analgesic and Sedative Effects

Most plant herbs have antipyretic and analgesic effects. The analgesic and sedative effects of agarwood can alleviate the clinical manifestations of gingivitis, such as swelling and pain. The agarwood contains a variety of analgesic and sedative substances, and valerenic acid has obvious sedative activity, α - Sandalyl alcohol and agarospirol have the same chlorpromazine like stabilizing effect as agallospirol. Through animal experiments, researchers found that agarwood oil has a sedative effect on

mice, which is manifested in that the autonomous motor activity of mice inhaled agarwood oil gradually decreases with the increase of medication time [28]. In addition, some scholars evaluated the pharmacodynamic effects of different parts of agarwood on sedation and hypnosis through the hypnosis experiment with pentobarbital sodium and the autonomous activity experiment. They found that the volatile oil of agarwood could significantly prolong the sleep time and increase the proportion of mice entering sleep. Based on the experimental results, the active molecules with sedative and hypnotic effects in the volatile oil of agarwood were further inferred [29]. In addition, they evaluated the sedative and hypnotic effects of agarwood essential oil through animal behavior experiments and explored the potential mechanism of action on GABAergic system. The results showed that agarwood volatile oil has a versatile sedative and hypnotic effect, and its mechanism of action may be related to regulating the gene expression of GABAA receptor, enhancing the function of GABAA receptor, and promoting the influx of Cl^{-} [30]. Researches have compared the analgesic effects of different years of agarwood, and deduce that the volatile oil of agarwood can significantly increase the pain threshold of mice and reduce the number of writhing induced by acetic acid in mice, proving that it has a clear analgesic effect [31]. The nonsteroidal anti-inflammatory drug diclofenac is a commonly used anti-inflammatory drug with analgesic and anti-inflammatory effects. The anti-inflammatory activity of agarwood essential oil in vivo and in vitro is comparable to that of standard diclofenac [9]. The analgesic and sedative effects of agarwood essential oil can be applied to gingivitis to alleviate the pain reaction caused by gingivitis and the active inflammation and immune reaction in the gums.

5. Immune Activity of Agarwood

The development of gingivitis is mostly due to the chronic inflammatory process caused by long-term microbial stimulation of the gums. Gingivitis involves a series of immune reactions, such as the accumulation of

inhibits the transformation of adaptive immune +t cells into Th1, Th2 and Th17 cells by inhibiting the differentiation of original CD4, and inhibits the activation, proliferation and differentiation of CD8+t cells and B cells. Inhibits innate and adaptive immunity by inhibiting STAT signaling pathway [33]. The excessive immunity of macrophages and neutrophils in inflammatory reaction is the main immune mechanism that causes gingivitis. Because the chromone and sesquiterpene in agarwood can inhibit the relevant signal pathway and immune response respectively, in conclusion, agarwood has the possibility to alleviate the overimmunity of gingivitis.

6. Conclusion

Sesquiterpenoids, flavonoids, quercetin, and other chemicals make up *Agaricus formosana*, the active ingredient in traditional Chinese medicine. There are a number of pharmacological effects shown by the active

lymphocytes, macrophages and plasma cells in the place where inflammation occurs. Overinduction, prolongation or imbalance of immune response will lead to immune diseases. Gingivitis is also an immune disease. Agarwood is not only able to enhance the cellular and humoral immune functions of the body, but also inhibit the immune function and reduce the release of inflammatory factors. This two-way immune regulation reflects the theory of "holism" and "balance between yin and yang" emphasized by traditional Chinese medicine [32]. Through modern pharmacological research, it has been found that agarwood and its essential oil have a variety of pharmacological activities, which may be related to their strong immune regulatory function. Inflammation, as the most basic immune reaction of the body, makes it possible to find new anti-inflammatory drugs from agarwood essential oil. Because of the correlation between immune reaction and inflammation, it was isolated from agarwood. Tryptone and sesquiterpene are not only the main anti-inflammatory substances, but also the main immunoactive substances. Compound CYF-2 is an epoxide 2 - (2-phenylethyl) - chromone derivative isolated from agarwood, which can significantly inhibit the activation of microglia, dendritic cells and neutrophils, which play an important role in innate immunity. In addition, GYF-21 significantly inhibits the transformation of adaptive immune-T cells into helper T cell 1 (Th1) and helper T cell 17 (Th17) by inhibiting the differentiation of primitive CD4, and inhibits the activation, proliferation and IFN of CD8- γ Secreting+t cells. Mechanism study showed that GYF-21 significantly inhibited STAT1/3 and NF- κ B in microglia- κ Activation of B signal path. In conclusion, GYF-21 can inhibit STAT1/3 and NF- κ B signal pathway significantly inhibits innate immunity and acquired immunity [7]. In addition, compound HHX-5 is a sesquiterpene derivative from agarwood. HHX-5 significantly inhibits the activation of macrophages and neutrophils, which plays an important role in innate immunity. In addition, HHX-5 strongly

components of agarwood, including anti-inflammatory, antioxidant, bacteriostatic, analgesic, sedative, and immunological properties. Plaque buildup on the gingival edge causes gingivitis, a persistent inflammatory condition. Finding a medication that can reduce inflammation without causing too many negative effects is a key area of focus in gingivitis research. In comparison to manufactured chemical medications, agarwood is safer. Nevertheless, agarwood is seldom used to treat gingivitis. By providing evidence of the aforementioned benefits, this study elucidates a potential mechanism by which agarwood reduces inflammation in periodontal tissues. For the agarwood industry to undergo a revolution, it offers a theoretical foundation.

References

- [1] [1] In an article published in 2008 in the Journal of Ethnopharmacology, the authors Zhou, Wang, Suolangjiba, and others investigated the anti-inflammatory and antinociceptive effects of an extract from the leaves of *Aquilaria sinensis* (Lour.) Gilg. Four Novel 2-(2-Phenylethyl) chromone Derivatives from Chinese Agarwood Derived via the Whole-Tree Agarwood-Inducing Method [2] Liu Y, Chen D, Wei J, et al. Published in 2016 in volume 21, issue 11, this article is referenced as 1433. the thirdThe authors of the article are Wang SL, Hwang TL, Chung MI, and others. Emerging Flavones, Anti-Inflammatory Components and a 2-(2-Phenylethyl)-4H-chromen-4-one Derivative Derived from *Aquilaria sinica* Stem Barks. The citation is from the journal *Molecules*, volume 20, issue 11, pages 20912–20925, published in 2015. Citation: [4] Liu Y, Chen H, Yang Y,... *Aquilaria sinensis* trees can be efficiently and innovatively produced high-quality agarwood using the whole-tree agarwood-inducing technology. *Chemical Communications*. 2013; 18(3): 3086-3106. This comes from the work of Shao H, Mei WL, Kong FD, and others [5]. Chinese agarwood "Qi-Nan" obtained from *Aquilaria sinensis* contains a novel 2-(2-phenylethyl) chromone glycoside. *Journal of Asian Natural Product Research*, 2017, 19, 1, 42–46. Agarwood alcohol extract inhibits oxidation and inflammation, protecting against gastric ulcers [6] (Wang C, Peng D, Liu Y, et al.). *Review of Alternative and Complementary Medicine*, 2021, 2021: 9944685. According to Guo R, Zhao YF, Li J, and others, GYF-21, a derivative of the epoxide 2-(2-phenethyl)-chromone, inhibits the STAT1/3 and NF- κ B signaling pathways, hence suppressing innate and adaptive immunity. Published in 2017 in the journal *Front Pharmacol*, volume 8, page 281. [8] Agarwood tea, a new modern herbal beverage made from the leaves of the *Aquilaria* (Thymelaeaceae) plant, and its pharmacological characteristics. Research by Adam AZ, Lee SY, and Mohamed R. Article published in 2017 in the *Journal of Herbal Medicine*, volume 10, pages 37–44. 9. Yadav DK, Mudgal V, Agrawal J, et al. Topical anti-inflammatory effectiveness of natural components of Agarwood oil studied by molecular docking and adverse drug event (ADEE) investigations. The current state of computer-assisted drug design is 360–370. Three novel sesquiterpenoids and one novel sesquiterpenoid derivative were isolated from Chinese eaglewood by Zhao, Peng, Han, and colleagues [10]. *Chemicals*. 2016; 21(3): 281. A study was conducted by Ma CT, Ly TL, Le THV, and colleagues on the anti-inflammatory effects of sesquiterpene derivatives derived from *Aquilaria malaccensis* agarwood on NO generation by macrophage RAW 264.7 cells [11]. *Journal of Phytochemistry*, 2021, 183: 112630. Inhibition of p-STAT3 and IL-1 β /IL-6 is the mechanism by which the essential oil of Chinese agarwood has an anti-inflammatory effect (Gao XL, et al., 2012). The article is published in the *Chinese Pharmaceutical Journal* in 2019 and has the DOI: 54/23/51-1957. This information is sourced from the work of R G Kumar L, Chatterjee NS, Tejpal CS, and others. Optimal chitosan wall material for spray-dried microencapsulation of squalene: oxidative stability experiments and characterization. *Worldwide Journal of Biomolecular Science*, 2017, 104 (Part B): 1986-1995. According to [14] Chen, Cai, Zhao, et al. A meta-analysis and comprehensive review of biomarkers in saliva and gingival crevicular fluid linked to oxidative stress and chronic periodontitis. *Journal of Clinical Periodontics*. 2019; 46(6): 608-622. A study conducted by Waddington, Moseley, and Embury [15] was conducted. Theories Behind Periodontal Disease: One possible component of periodontal disease pathophysiology is reactive oxygen species. *Journal of oral disorders*, 2000, 6, 3, 138–151. [16] GC-MS by Wang, Li, Luo, and colleagues An analysis of the chemical constituents of agarwood and several *Aquilaria sinensis* (Lour.) Gilgorgans from various Asian nations. The article is published in *Molecules* on April 23, 2018, volume 23, issue 9, pages 2168. [17] In a study conducted by Dahham, Tabana, and Iqbal et al. The antibacterial, anticancer, and antioxidant characteristics of the sesquiterpene β -caryophyllene derived from the *Aquilaria crassna* essential oil. Citation: *Molecules*, 2015, 20(7): 11808-11829. The essential oil of *Lignum Aquilariae Resinatum* protects PC12 cells from oxidative damage caused by hydrogen peroxide, according to Xiong LY et al. [18]. *Chinese Journal of Pharmaceutical Research*, Volume 25, Issue 1, Pages 28–32, 2014. By Parwata, Manuaba, and Yasa [19] we mean... *Gyrinops versteegii* leaves, an extract of which is water-based, contain powerful flavonoid components that function as antioxidants. *Journal of Biomedical and Pharmacology*, 2018, 11(3): 1501-1511. According to Ma, Qiao, Fu, et al. (2021). The Biological Activity of Man-Made and Naturally Occurring Agarwood: A Comparative Study. Article number: 1532 in *Forests*, volume 12, issue 11. Reference: [21] Batubara R, Wirjosentono B, Siregar AH, and colleagues. Antimicrobial action

against skin-dwelling bacteria and fungus, as well as bioactive components of an ethanol extract of agarwood leaves (*Aquilaria malaccensis*). Publication: Biodiversitas, Volume 22, Issue 1, Pages 2884–2890, 2021.

Mikaido H. [22] Bacterial cell wall permeability. "Angew Chem International, English Edition," 1979, 18, 5, 337–350.

Proteinase inhibitor gene families: methods for enhancing herbivore resistance via genetic engineering [23] Ryan CA. Volume 10, Issue 1, pages 20–24, 1989, Bioessays.

[24] Huang Y, Wu J, Zeng Y, et al. The chemical make-up and anti-MRSA efficacy of Chinese eaglewood essential oil. The citation is from the Journal of Chinese Pharmaceutical Science, volume 17, issue 3, page 225.

Essential oil and heartwood extracts of *Aquilaria crassna* produced by water distillation and supercritical fluid carbon dioxide extraction: chemical components and antibacterial activity [25] Wetwitayaklung P, Thavanapong N, Charoenteeraboon J. Volume 2009, Issue 3, Pages 25–33, Journal of Science, Technology, Engineering, and Health Affairs.

Cushnie and Lamb [26] said. A better knowledge of flavonoids' antimicrobial activities has recently emerged. Journal of antimicrobial agents: an international publication, 2011, 38 (2), 99-107.

This sentence is paraphrased from a publication by Takemoto et al. [27]. Exploring the sedative effects of vaporizing spikenard extract and agarwood oil, as well as identifying their active ingredients. "Journal of the National Medical Association" (2008), 62(1), 41–46.

Wang S. et al. [28] investigated the effects of agarwood made using the whole-tree agarwood-inducing approach on the suppression of spontaneous activity and hypnosis in mice. Published in the Journal of International Pharmaceutical Research in 2016, volume 43, pages 1082–1089.

An Essential Oil from Agarwood Demonstrates GABAergic-Mediated Sedative-Hypnotic Effects [29] Wang et al. "Molecules" published in 2017 with the DOI 22/12: 2190.

[30] Research by Wang et al. comparing the analgesic effects of Chinese eaglewood from various years. (Hainan Medical Journal, 2014, 25: 2188-2190).

The article "Aquilaria Species (Thymelaeaceae) Distribution, Volatile and Non-Volatile Phytochemicals, Pharmacological Uses, Agarwood Grading System, and Induction Methods" was written by Wang Y, Hussain M, Jiang Z, and others—[31]. Journal of Molecules, 2021, 26, 24: 7708.

The authors of the piece are Ma HD, Deng YR, Tian Z, and Lian ZX. The relationship between

TCM and immunological modulation. Clinical Review of Allergy and Immunology. 2013; 44(3): 229–241.

According to research by Zhu, Zhao, Huo, and colleagues, HHX-5, a sesquiterpene derivative found in Chinese agarwood, inhibits STAT signaling pathways and hence reduces both innate and adaptive immunity [33]. European Journal of Pharmacology, 2016, 791: 412-423.